
Patient Name

Physician

Procedure Date/Arrival Time

Procedure Time

Location:

- McLaren Hospital - Pennsylvania**
2727 S. Pennsylvania Ave, Lansing
- McLaren Hospital- Greenlawn**
401 W. Greenlawn Ave, Lansing
- Sparrow Hospital**
1215 E. Michigan Ave, Lansing
- Sparrow Health Center**
2909 E Grand River, Lansing
- Sparrow St Lawrence**
1210 West Saginaw Street, Lansing
- Lansing Surgery Center**
1707 Lake Lansing Rd, Lansing
- Hayes Green Beach Hospital**
321 E. Harris St, Charlotte
- Eaton Rapids Specialty Clinic**
1500 S. Main St. Eaton Rapids

As of 02/1/2020 MGI & CCS will implement a rescheduling fee for any rescheduling of a procedure more than 3 days after being booked.

1st Rescheduled appointment is \$25

2nd Rescheduled appointment is \$50

3rd Rescheduled appointment is \$75

If you cancel your procedure for any reason within 5 business days of the scheduled procedure, a \$100 fee will be assessed to your account

Results and Follow Up

- Please go home and rest for the remainder of the day. Do not drive or work or operate machinery for the rest of the day.
- Your physician will discuss test results with you and your family member following the procedure.
- You will be given written instructions for diet, activity, and follow up instructions.
- If biopsies were taken, a results letter will be sent through our patient portal. If not a member of Follow My Health, the letter will be mailed to you within 14 days of your procedure.

Important Driving Instructions

You must have an adult 18 years or older remain with you during the procedure and drive you home afterward. Your procedure will be cancelled if you fail to meet these requirements.

Billing Procedure

There may be up to four charges associated with your procedure (Physician, Hospital/Facility, Anesthesia, and Lab). Please verify with your insurance carrier your benefit coverage for each.

Please be sure you bring your picture ID and insurance cards with you to the facility the day of the procedure. Failure to do so will result in cancellation of your procedure.

Digestive Health Institute



MIRALAX EXTENDED PREPARATION

Board Certified Gastroenterologists

Iftiker Ahmad, M.D.
Oussama Al Sawas, M.D.
Radoslav Coleski, M.D.
Dorian Jones, M.D.
Scott Plaehn, D.O.
Robert Rose, D.O.
Albert Ross, M.D.
Dana Stewart, D.O.
John Walling Jr., D.O.
David Wiedemer, M.D.
Siaka Yusuf, M.D.

Board Certified Colorectal Surgeons

Daniel C. Coffey, M.D.
Lucas Julien, M.D.
Razvan Opreanu, M.D.

If you have questions, please contact _____ at (517) 332-1200 ext _____

For more information visit our website www.michigangastro.com
www.capitolcolon.com

Colonoscopy

The purpose of this procedure is to directly visualize the mucosal lining of the colon in order to inspect for disease. The principal risk of colonoscopy is the remote possibility of perforation. Immediately prior to the passage of the instrument, you may be given intravenous Demerol, Versed, or Propofol. If you are allergic to any of these, latex or eggs please be certain you have notified the scheduler.

Medication

If you are taking blood thinners, Aggrenox, Plavix, Pradaxa, Effient, Lovenox, Ticlid, Coumadin, Trental, Persantine, Heparin, or Fragmin) please be certain you have notified the scheduler.

You must discontinue any diet medication 1 week prior to your procedure

Labs

Please have any ordered bloodwork drawn

_____ days before your test. A lab requisition for your bloodwork has been enclosed.

_____ You do not need bloodwork at this time

NOTE: Female patients of menstruating years will be required to provide a urine specimen when you arrive to the hospital/facility.

Five Days Prior To Procedure:

Please refrain from eating foods that contain nuts, seeds or corn. You will also need to stop taking fiber and iron supplements.

Two Days Prior To Procedure:

1. **CLEAR LIQUID DIET all day long** (this starts at 12:00am or when you wake up in the morning. **No solid foods**. Drink an extra 8-oz of clear liquids every hour throughout the entire day.

2. **Diabetics:** Use ½ of your usual insulin dose. Do not take your diabetic pills today.

3. At 7 PM drink one bottle of Magnesium Citrate.

4. Mix entire contents of Miralax with 64 oz (2 liters) of clear liquids in a tall pitcher. Mix well and refrigerate. Mixture will be used tomorrow and the day of procedure.

Day Before Examination

1. **Continue clear liquid diet all day. NO SOLID FOOD.** Drink 8-oz of clear liquids hourly.

2. **Diabetics:** Use ½ of your usual insulin dose. Do not take your diabetic pills today.

3. At 3:00 PM take 1 dulcolax tablet and continue drinking clear liquids.

4. At 6:00 PM begin drinking 1 cup (8 oz) of the of the laxative solution every 15 minutes (total of 4 cups) You must finish drinking the solution within 1 hour (**32 oz. total**) to ensure a clean colon. It is not unusual to have nausea and vomiting from this medication. If this occurs, stop the laxative, suck on mints or hard candy and resume the drinking the laxative 1 hour later. You should continue clear liquids 8 oz hourly throughout the evening. If there is no bowel movement or stools are not turning clear by 10 pm, please drink 1 bottle of Magnesium Citrate.

Day of Examination

1. You may only have clear liquids the day of your test (no Ensure this day) to complete your prep.

2. Six hours before the start of your test begin drinking 1 cup (8 oz) of the laxative solution every 15 minutes (total of 4 cups). You must finish drinking the solution within 1 ½ hours (**32 oz. total**) then **NOTHING else by mouth prior to your procedure.** You must be finished with your prep at least 4 1/2 hours prior to your procedure then nothing by mouth. (No water, gum, mints, hard candy, Ensure, chewing tobacco etc) until after the procedure is completed. Failure to do so will result in cancellation of your procedure(s).

3. **Diabetics:** Please check your blood sugar and take this information with you to the hospital/facility. **Do not** take your insulin or diabetic pills this morning. Bring all of your insulin with you to the procedure.

4. Please only take your heart, blood pressure, seizure, respiratory, anti-rejection or anti-anxiety medications at 6 AM with a small sip of water; you may use your inhalers.

Clear Liquid Diet

- Coffee, tea, or cola
- Apple, white grape, or white cranberry juice
- Up to 3 cans or bottles of vanilla or butterpecan Ensure or Glucerna (diabetics) that are labeled "suitable for lactose intolerant".
- Plain jello (no red colors)
- Clear soups and/or broth (strain off all vegetables and/or noodles)
- Popsicles (no red colors)
- Artificially sweetened powdered drinks (kool-aid, tang, crystal light – no red or purple colors)
- Sorbet that does not contain milk or chunks of fruit
- **No** milk or milk byproducts (cheese, yogurt)
- **No** grapefruit, tomato, V-8, or orange juice
- **No** alcohol

Shopping List:

- One bottle of MiraLax (238-grams)
- One dulcolax or bisacodyl laxative tablet (not stool softener)
- One bottle Magnesium Citrate (10 oz)
- Two 32-oz bottles of Powerade-lemon-lime **or** four 16 oz bottles of Propel, **or** any 64 oz (2 liters) any (non-red/purple) clear liquid. *This will be used to mix with the Miralax.*
- Hard candy or peppermints if you become nauseated.